

DNA Methylation Screening Identifies Driver Epigenetic Events of Cancer Cell Survival

Daniel D. De Carvalho,^{1,3} Shikhar Sharma,^{1,2,3} Jueng Soo You,¹ Sheng-Fang Su,^{1,2} Phillippa C. Taberlay,¹ Theresa K. Kelly,¹ Xiaojing Yang,¹ Gangning Liang,¹ and Peter A. Jones^{1,*}

¹Department of Urology, Biochemistry and Molecular Biology

²Program in Genetic, Molecular and Cellular Biology

University of Southern California/Norris Comprehensive Cancer Center Keck School of Medicine, University of Southern California, Los Angeles, CA 90089-9181, USA

³These authors contributed equally to this work

*Correspondence: pjones@med.usc.edu

DOI 10.1016/j.ccr.2012.03.045

SUMMARY

Cancer cells typically exhibit aberrant DNA methylation patterns that can drive malignant transformation. Whether cancer cells are dependent on these abnormal epigenetic modifications remains elusive. We used experimental and bioinformatic approaches to unveil genomic regions that require DNA methylation for survival of cancer cells. First, we surveyed the residual DNA methylation profiles in cancer cells with highly impaired DNA methyltransferases. Then, we clustered these profiles according to their DNA methylation status in primary normal and tumor tissues. Finally, we used gene expression meta-analysis to identify regions that are dependent on DNA methylation-mediated gene silencing. We further showed experimentally that these genes must be silenced by DNA methylation for cancer cell survival, suggesting these are key epigenetic events associated with tumorigenesis.

INTRODUCTION

During tumorigenesis cancer cells acquire, through a multistep process, a new set of properties that allows them to overcome physiological homeostasis. These properties include unlimited proliferation potential, self-sufficiency in growth signals, resistance to antiproliferative and apoptotic signals and immune system evasion, among others (Hanahan and Weinberg, 2000, 2011). These alterations, on the other hand, contribute to a process known as the stress phenotype of cancer (Luo et al., 2009), which includes DNA damage/replication stress, proteotoxic stress, mitotic stress, metabolic stress, and oxidative stress.

To survive the tumorigenic process, a cancer cell undergoes several modifications to its genomic circuitry, such as activating mutations in oncogenes and aberrant activation of nononcogenic pathways. These adaptations lead to oncogene addiction (Weinstein, 2002) and nononcogene addiction (Solimini et al.,

2007), respectively. Because of this aberrant circuitry, cancer cells become hypersensitive to the effects of classic tumor suppressor genes (TSGs) (Luo et al., 2009; Weinstein, 2002) and, potentially, to genes that can inhibit the nononcogenic signaling pathways that cancer cells rely on to survive.

Changes in the cancer cell transcriptome can be driven by genetic and epigenetic alterations (Baylin and Ohm, 2006; Jones and Baylin, 2007). DNA methylation is an epigenetic process that can heritably change gene expression without altering the DNA sequence. In normal somatic cells, most DNA methylation occurs at CpG dinucleotides within CpG poor sequences, whereas CpG-rich sequences, also known as CpG islands, are usually unmethylated (Sharma et al., 2010). DNA methylation is a vital mechanism of epigenetic gene silencing, playing key roles in X chromosome inactivation, genomic imprinting, embryonic development, silencing of repetitive elements and germ cell-specific genes, differentiation, and maintenance of pluripotency (De Carvalho et al., 2010; Meissner, 2010; Robertson,

Significance

Epigenetic modifications are potentially reversible, making them good “druggable” targets. Here, we show that cancer cells cannot survive in the absence of aberrant DNA methylation of specific promoter regions. This process may render cancer cells more susceptible to epigenetic therapy. We also found that physiological DNA methylation of germline-specific genes is necessary for somatic cell survival, suggesting a physiological dependence on continuous DNA methylation of these regions in somatic tissues. Moreover, by defining the promoter regions that must be methylated in order for cells to survive in culture, we found several genes that acquire de novo DNA methylation in cell lines, highlighting the importance of careful interpretation of epigenetic results obtained from cell culture experiments.




Figure 1. Clustering of DNMT-Deficient Cells Identifies Three Classes of Putative Driver Genes Marked by DNA Methylation

(A) One-dimensional hierarchical clustering using Euclidean distance and average linkage was performed with the ~24,000 Infinium DNA methylation probes located outside of repeats or known SNPs in HCT116 wild-type, DKO8, and DKO1 cell lines. Each row represents a probe; each column represents a sample. The β value (level of DNA methylation) for each probe is represented with a color scale as shown in the key.

(B) k Means ($K = 4$) clustering of the 566 Infinium DNA methylation probes that maintain DNA methylation in DKO1 sample (a β value of at least 0.6 and a difference between HCT116 and DKO1 smaller than 0.2) in (A) for 10 TCGA samples ($n = 4$ normal colon and $n = 6$ primary colon adenocarcinoma).

(C) Heat map of 566 Infinium DNA methylation probes in 32 normal tissues retaining the probe order from (B). Primary normal bladder ($n = 4$), sperm ($n = 1$), and primary normal TCGA kidney ($n = 15$), lung ($n = 4$), and ovary ($n = 8$). WGA DNA was used as a negative control for DNA methylation.

(D) Bisulfite-sequencing validation of Infinium DNA methylation data from two regions (*LDHAL6B* and *ADAM2*) from the somatic-specific DNA methylation cluster and two regions (*ARMCX1* and *MEOX2*) from the cancer-specific DNA methylation cluster. Arrow indicates the position of the Infinium probe. Empty and filled

2005). Besides these physiological roles, deregulated DNA methylation can also be a major driver of pathological conditions, including neurological and autoimmune diseases, as well as cancer (Kelly et al., 2010; Portela and Esteller, 2010; Taberlay and Jones, 2011). During tumorigenesis, global DNA methylation patterns change, resulting in hypomethylation of non-CpG islands and hypermethylation of CpG islands (Sharma et al., 2010). DNA hypermethylation has been shown to result in abnormal silencing of several TSGs in most types of cancer (Jones and Baylin, 2002, 2007).

Recently, several efforts to examine the cancer methylome, utilizing genome-wide techniques, have revealed that a large number of genes exhibit aberrant DNA methylation profiles in cancer (Figuerola et al., 2010; Irizarry et al., 2009). These changes can be used to stratify subtypes of cancers (Figuerola et al., 2010; Noushmehr et al., 2010) and to predict cancer outcomes (Portela and Esteller, 2010), among other uses. Distinguishing which genes play key “driver” roles via DNA methylation-mediated gene silencing in cancer initiation, progression, and maintenance and those genes that are only “passengers” in the tumorigenic process would be extremely useful in developing more targeted epigenetic therapies (Kelly et al., 2010). However, making this distinction has proven extremely difficult due to the large number of differentially DNA-methylated genes in human cancers (Kalari and Pfeifer, 2010).

We, and others, have suggested that cancer cells may become addicted to an aberrant epigenetic landscape, especially with respect to DNA methylation (Baylin and Ohm, 2006; Kelly et al., 2010). However, as of yet, and to our knowledge, there is no direct evidence for such an addiction. Furthermore, mining the thousands of genomic regions that are de novo DNA methylated in cancer and identifying those required for cancer cell survival have proven extremely challenging (Kalari and Pfeifer, 2010). Here, we describe an approach to identify driver epigenetic events associated with cancer cell survival. Our findings pave the way for new generations of epigenetic therapies, which target the genes cancer cells rely on being silenced by DNA methylation in order to survive.

RESULTS

Identification of the Minimum DNA Methylation Profile Required for Cancer Cell Survival

We hypothesized that cancer cells depend on DNA methylation of a few key regions for survival and that these regions would preferentially maintain methylation when artificially reducing global DNA methylation. To test this hypothesis, we profiled HCT116 colon cancer cells and HCT116 cells with a genetic disruption of *DNMT3B* and *DNMT1* (DKO) (Rhee et al., 2002). This genetic disruption led to a complete knockout of *DNMT3B* and a truncated *DNMT1* transcript, expressed at very low levels (Egger et al., 2006; Rhee et al., 2002; Spada et al., 2007). For this study we used two DKO subclones, DKO8 and DKO1, which

retain approximately 45% and 5% of the HCT116 wild-type global DNA methylation levels, respectively (Rhee et al., 2002; Sharma et al., 2011). It is important to note that a further reduction of *DNMT1* levels, by RNAi, in cells with a genetic disruption of *DNMT1* results in demethylation and a massive reduction of cell viability and immediate induction of cell death (Spada et al., 2007), suggesting that DNA methylation is required for cancer cell survival.

We profiled promoter DNA methylation of HCT116, DKO8, and DKO1 cell lines using the Illumina Infinium platform (HumanMethylation27) and observed a reduction in global DNA methylation levels in DKO8 cells compared to HCT116 wild-type cells and an even greater reduction in DKO1 cells (Figure 1A), consistent with previous data (Rhee et al., 2002; Sharma et al., 2011). Surprisingly, we found a collection of 566 CpG sites, spanning 490 genes that despite the strong impairment in DNA methyltransferase activity, still retained a high level of DNA methylation in DKO1 cells, with a β value higher than 0.6 (see Table S1 available online for gene/probe list). These regions were also highly methylated in HCT116 and DKO8 cells, and none showed a difference in their β values greater than 0.2 among the three cell lines.

Next, we sought to identify whether there was a cancer-specific DNA methylation profile at these regions that maintained DNA methylation even in DKO1 cells, which would potentially include important putative targets for epigenetic therapy. To accomplish this, we first compared the DNA methylation levels of the 566 CpG sites that retained DNA methylation in DKO1 cells to the DNA methylation profile of 6 primary colon adenocarcinoma tissue samples and 4 normal colon tissue samples obtained from The Cancer Genome Atlas (TCGA) database. Using k means clustering, we identified 92 CpG sites, spanning 77 genes that were unmethylated in normal colon and became hypermethylated in colon adenocarcinoma (Figure 1B; Table S1), consistent with a cancer-specific methylation profile.

We further compared these data to DNA methylation data of several normal tissues including sperm, bladder, kidney, lung, and ovary, which allowed us to identify clusters of gene regions highly enriched for somatic tissue-specific DNA methylation. Such genes were methylated in the somatic tissues analyzed and unmethylated in germ cells. This somatic tissue-specific cluster comprised 99 CpG sites, spanning 83 genes (Figure 1C; Table S1). Furthermore, we also identified genes that exhibit cell culture-specific DNA methylation, such that these regions are methylated in all cell lines analyzed but unmethylated in primary tissues (Figure 1C; Table S1). This cell culture-specific cluster comprised 29 CpG sites, spanning 25 genes. We focused only on these three groups because of their differential DNA methylation profiles. We speculate that the remaining 346 CpG sites might be regions that are more prone to methylation, remaining a good target for residual DNA methylation activity without functional relevance or, alternatively, may have a tissue-specific expression profile, being unmethylated only in

circles denote unmethylated and methylated CpG sites, respectively. Each horizontal row represents one sequenced DNA clone. The number on the right represents the mean DNA methylation score of each region, and the number in the parentheses represents the mean DNA methylation score of the specific Infinium CpG site.

See also Figure S1 and Table S1.

specific cell types that were not surveyed in this study. Whole Genome Amplified (WGA) DNA served as a negative control (Figure 1C) to confirm that the regions identified as being methylated in DKO1 cells were not false positives due to technical problems with the specific Infinium probes.

The distribution of probes, relative to transcription start sites (TSSs), in the cancer-specific and somatic tissue-specific clusters was found to be very similar to the distribution of the array itself, whereas the distribution in the cell culture-specific cluster tended to be slightly more concentrated at the TSS (Figure S1A). However, it should be noted that there was no association between distance to TSS and methylation cluster (somatic, cancer, cell line) as assessed by one-way ANOVA ($p > 0.05$).

We selected genomic regions from each cluster to validate the Infinium-based DNA methylation data using bisulfite sequencing. All of the sequences analyzed showed high levels of DNA methylation in HCT116 wild-type and DKO1 cells (Figure 1D), with the CpG site surveyed by the Infinium platform presenting a maximum difference between their β values of 0.17 in DKO1 when compared to HCT116 wild-type. These results demonstrate that even though DKO cells are globally DNA hypomethylated (Rhee et al., 2002; Sharma et al., 2011), the residual DNA methylation is focal and site specific, supporting the hypothesis that there is a functional role for some of the retained DNA methylation.

To further demonstrate the importance of the DNA methylation that is retained in the three identified clusters, we treated DKO1 cells with 1 μ M of 5-Aza-2'-deoxycytidine or PBS for 24 hr. After treatment, we allowed at least two population doublings (5 days) for demethylation to occur and then analyzed changes in the DNA methylation profile by the Illumina Infinium array. As expected, most of the regions in the three previously identified clusters were resistant to demethylation, with only eight regions from the somatic cluster, five regions from the cancer cluster, and one region from the cell culture cluster presenting a difference in the β value greater than 0.2 (Figure S1B). These regions we considered false positives and excluded them from subsequent analysis.

Residual Methylation in DKO1 Cannot Be Explained by an Inherent Susceptibility to DNA Methylation

Our working hypothesis is that the artificial impairment of DNA methyltransferase machinery in DKO1 cells will induce a strong selective pressure for any remaining DNA methylation to be maintained at the regions necessary for cancer cell survival. An equally plausible hypothesis is that the residual methylation reflects an inherent tendency for some genes to remain methylated. Indeed, previous studies suggest that certain genomic regions are more prone to DNA methylation (Estéicio et al., 2010; Ohm et al., 2007; Schlesinger et al., 2007; Widschwendter et al., 2007). Therefore, these regions may remain better targets for residual DNA methylation activity. To directly test this alternative hypothesis, we used two known approaches to predict whether a gene is more prone to DNA methyltransferase activity in cancer cells: one based on its chromatin structure (Ohm et al., 2007; Schlesinger et al., 2007; Widschwendter et al., 2007), and another based on its genomic architecture (Estéicio et al., 2010).

Genes marked by H3K27me3 in embryonic stem cells (ESCs) are known to be predisposed to DNA methylation in cancer cells (Ohm et al., 2007; Schlesinger et al., 2007; Widschwendter et al., 2007). Indeed, we found that the H3K27me3 status in ESCs can accurately predict the methylation levels in the wild-type HCT116 cells (Figure 2A). Then, we tested whether methylation-prone regions (H3K27me3 positive in ESCs) would preferentially retain methylation in DKO1 cells, compared to HCT116 cells. If this hypothesis was correct, we should observe an enrichment of methylation-prone genes in the cohort of genes that is methylated in DKO1 cells because they would be better targets for residual DNA methyltransferase activity. Yet, there was no such enrichment (Figure 2B). Rather, we observed a slight decrease in the proportion of methylation-prone genes that retain methylation in DKO1 cells. These data suggest that genes found to retain DNA methylation in DKO1 cells are not simply predisposed to DNA methylation in cancer cells.

To further test this hypothesis, we performed a similar analysis using a previously published algorithm to predict whether a genomic region is prone, intermediate, or resistant to DNA methylation in cancer cells, based on its genomic architecture (Estéicio et al., 2010). In our test this algorithm accurately predicted methylation levels in HCT116 cells (Figure 2C). Similar to the previous analysis, if the genes that maintain methylation in DKO1 cells were simply more prone to DNA methyltransferase activity, one would expect an enrichment of methylation-prone genes in the pool of genes that retains methylation in DKO1 cells. Again, we could not find such enrichment (Figure 2D).

Taken together, these data suggest that the targets of residual DNA methylation in DKO1 cells are not dictated by an inherent predisposition to DNA methylation based on either the chromatin structure or the genomic architecture. These findings further support our original hypothesis that these loci retain methylation due to a functional selection pressure.

Validation of the Findings in Other Types of Cancer and Association with Gene Expression

We next validated our findings in a larger test set of colon adenocarcinoma and normal colon samples. Using DNA methylation data available from TCGA, we observed a significant increase in DNA methylation in the majority of the CpG sites identified with cancer-specific DNA methylation in 168 primary colon adenocarcinoma samples relative to 16 normal colon samples (Figure 3A; Wilcoxon rank sum test followed by a FDR correction, $p < 0.05$). We next extended our findings to determine whether this cancer-specific DNA methylation profile was unique to colon adenocarcinoma or if it could also be observed in other tumor types. Using DNA methylation data available from TCGA, we analyzed DNA methylation from 19 lung adenocarcinoma samples against 4 normal samples. Again, the same pattern emerged, where the identified CpG sites presented an overall significant gain of DNA methylation in the tumor samples (Figure 3C; Wilcoxon rank sum test followed by a FDR correction, $p < 0.05$). Indeed, most of the genes statistically determined to be hypermethylated in lung adenocarcinoma were also hypermethylated in colon adenocarcinoma (Figure 3E). These data indicate that the CpG sites we identified as cancer specific are frequently hypermethylated in other types of human cancer relative to the normal cell counterparts, suggesting that these




Figure 2. Residual Methylation in DKO1 Is Not Caused by an Inherent Susceptibility to DNA Methylation

(A) Validation of H3K27me3 status in ESCs as a predictive method for DNA methylation in HCT116 cells. Methylation status of ~27,000 CpG sites was determined by Infinium. A t test with Mann-Whitney U posttest was performed. Data represent the mean \pm SEM.

(B) Frequency of probes marked by H3K27me3 in ESCs in the cohort of DNA-methylated probes (β value > 0.6) in HCT116, DKO8, and DKO1 cells.

(C) Validation of the predictive method based on genomic architecture (Estéicio et al., 2010) in HCT116 cells. Methylation status of ~27,000 CpG sites was determined by Infinium. One-way ANOVA with Kruskal-Wallis test as performed. Data represent the mean \pm SEM.

(D) Frequency of methylation-prone genes in the cohort of DNA-methylated genes (β value > 0.6) in HCT116, DKO8, and DKO1 cells.

regions might have a more fundamental role in tumorigenesis, such as cell survival.

Because DNA methylation of CpG islands located in promoter regions is well known to be correlated with gene silencing (Cedar and Bergman, 2009; Jones and Baylin, 2007; Portela and Esteller, 2010), we investigated the expression state of the genes identified using independent data sets. We selected two microarray data sets from the Gene Expression Omnibus database (<http://www.ncbi.nlm.nih.gov/geo/>): colon adenocarcinoma against normal colon (GSE 8671) (Sabates-Bellver et al., 2007); and lung adenocarcinoma against normal lung (GSE7670) (Su et al., 2007).

We found an inverse correlation between DNA methylation and gene expression when we analyzed the gene expression data of 32 normal colon samples and 25 colon tumor samples. The majority of the genes subject to cancer-specific DNA methylation displayed decreased gene expression in colon cancer samples compared to normal colon (Figure 3B; t test followed by a FDR correction, $p < 0.05$). We also observed that some genes showed a similarly low level of expression in both samples, probably due to an epigenetic switch in the silencing mechanism where the gene was already silenced in the normal sample by another epigenetic mechanism and became de novo DNA methylated in cancer (Gal-Yam et al., 2008).

Moreover, we found a similar gene expression pattern in lung adenocarcinoma, where most of the cancer-specific DNA

methylation genes displayed decreased gene expression in the tumor when compared to the correspondent normal tissue (Figure 3D; t test followed by a FDR correction, $p < 0.05$). Again, most of the genes statistically repressed in lung adenocarcinoma were also repressed in colon adenocarcinoma (Figure 3F). These data further suggest that there is a functional relevance of identified DNA methylation.

Altogether, by combining gene expression with DNA methylation data, we identified regions that are candidates for DNA methylation-mediated gene silencing. Moreover, the gene expression data corroborate our cluster analysis by using a different method and independent data sets to demonstrate biological differences in the gene clusters we identified.

Spontaneous Loss of DNA Methylation at the Identified Genomic Regions Is Associated with Cell Death

DKO1 cells have highly impaired DNA methyltransferase machinery due to the absence of DNMT3B, very low protein levels of DNMT3A, and low levels of a truncated DNMT1 (Egger et al., 2006; Sharma et al., 2011). As a consequence of the impaired DNA methyltransferase machinery, the global DNA methylation level in this cell line is very low, with most of the genes that were methylated in the parental HCT116 cells losing this methylation in DKO1 cells. Therefore, we hypothesize that DKO1 cells would be under a constant selective pressure to maintain the residual DNA methylation at key regions necessary for this cancer cell to survive.




Figure 3. Validation of CpG Sites Identified with Cancer-Specific DNA Methylation Using Independent Data Sets and Association with Gene Repression

(A) Volcano plot of the CpG loci identified as cancer-specifically methylated in colon adenocarcinoma (normal n = 16, cancer n = 168) from TCGA Data Portal. The β value difference in DNA methylation between the tumor samples and the correspondent normal samples is plotted on the x axis, and the p value for a FDR-corrected Wilcoxon rank sum test of differences between the tumor and correspondent normal samples ($-1 \times \log_{10}$ scale) is plotted on the y axis. Probes that are significantly hypermethylated (FDR adjusted $p < 0.05$) in tumors are shown in red.

(B) Volcano plot gene expression data of cancer-specific DNA-methylated genes. Gene expression data were obtained from GEO (GSE 8671) from primary normal colon (n = 32) and primary colon cancer (n = 25). For the volcano plots, gene expression fold change between the normal tissues and the tumor tissues is

We next investigated whether DKO1 cells exhibit a higher basal level of cell death than HCT116 wild-type cells. When quantifying cell death by measuring the externalization of phosphatidylserine (PS) using annexin V by flow cytometry, we observed at least four times more spontaneous cell death in DKO1 than in the parental HCT116 cells (Figure 4A). This suggests that DKO1 cells are indeed under constant selective pressure, probably because during cell division, some daughter cells lose DNA methylation at key regions due to the impaired DNA methyltransferase activity in DKO1 cells (Egger et al., 2006; Spada et al., 2007), and consequently, they cannot survive.

We took advantage of the increased rates of spontaneous cell death in DKO1 cells to further test our hypothesis that cancer cells depend on constant DNA methylation of these regions in order to survive. Using cell sorting, we first separated DKO1 cells into two populations: annexin V positive (early spontaneous apoptosis), and annexin V negative (viable cells) (Figure 4A). These two populations have distinct morphologies, with annexin V-positive cells in the range of lower Forward Scatter (FSC) and higher Side Scatter (SSC), a characteristic feature of apoptotic cells (Darzynkiewicz et al., 1992), compared to annexin V-negative cells (Figure 4B).

We then compared the DNA methylation levels of *EYA4* and *IRAK3* gene promoter regions in early apoptotic and viable cells. We have previously defined these genes as harboring cancer-specific DNA methylation and differential expression in cancer versus normal cells. Furthermore, these genes were in the top tier for significantly hypermethylated genes, and for gene repression, in colon and lung adenocarcinoma. In addition we also compared the DNA methylation levels of *SYCP3* and *ADAM2* gene promoter regions between early apoptotic and viable cells. These genes were identified as having somatic cell-specific DNA methylation and differential gene expression between somatic and germ cells (data not shown). In agreement with our hypothesis that DNA methylation-induced silencing of these regions is required for survival, early apoptotic cells showed at least a 27% reduction in DNA methylation in all four regions analyzed, with some specific CpG sites having as much as 80% reduction in DNA methylation (Figures 4C and S2A). Because degradation of cellular mRNA is an early apoptosis-induced event (Del Prete et al., 2002), we could not reliably measure whether this demethylation was associated with re-expression of these genes in the dying cells. In contrast, DNA degradation is a late apoptotic event, which allowed us to study the DNA methylation status during the first steps of apoptosis.

An alternative hypothesis is that global demethylation in DKO1 cells, due to impaired DNA methyltransferase activity, leads to genomic instability and cell death. To test this hypothesis, we measured the global DNA methylation levels of the early spontaneous apoptotic and viable cells and did not find a global reduction in DNA methylation (Figure S2B), further suggesting that demethylation of these specific genes lead to cell death. In addition to test whether apoptosis itself could cause demethylation of these regions, we treated HCT116 cells with 0.2 μ M of Staurosporine (STS), a drug known to induce cell death by blocking protein kinases (Manns et al., 2011). Next, we sorted viable and STS-induced dead cells and did not observe any difference in DNA methylation of these candidate regions (Figure S2C). Altogether, this strongly suggests that demethylation of these regions is causing cell death rather than the other way around.

These data, together with our previous data showing that DKO1 cells have reduced cell viability when this low level of DNMT1, and consequently the DNA methylation level, is further reduced by RNAi (Egger et al., 2006; Spada et al., 2007), and that complete knockout of the maintenance *DNMT1* leads to massive cell death (Chen et al., 2007), demonstrate that these cells are under constant selective pressure to retain DNA methylation at these key regions that we identified here in order to survive.

Functional Validation

We sought to further demonstrate that re-expression of genes whose DNA methylation is critical for cancer cell survival leads to increased cell death. We cloned the cDNA of six genes from the cancer cluster (*IRAK3*, *P2RY14*, *CDO1*, *BCHE*, *ESX1*, and *ARMCX1*), two from the somatic cluster (*ADAM2* and *SYCP3*), and one from the cell line cluster (*STEAP4*) into the pLJM1 lentiviral vector to individually reexpress these genes in HCT116 and RKO colon carcinoma cell lines (Figures S3A and S3B). We observed that expression of each of these genes decreased cell viability in both HCT116 and RKO cells (Figures 5A and 5B). We also re-expressed *NOX4* as a control gene (Figures 5A and 5B). *NOX4* was heavily methylated in HCT116 (β value of 0.95) and completely demethylated in DKO1 (β value of 0.007), suggesting that DNA methylation-mediated repression of this gene is not necessary for DKO1 survival. It is important to note that these ten genes have a low relative expression in RKO (Figure S3B) and a very high basal DNA methylation level in this cell line (Figure S3E).

plotted on the x axis, and the p value for a FDR-corrected t test of differences between the normal and the tumor tissues ($-1 \times \log_{10}$ scale) is plotted on the y axis. Probes that are significantly ($p < 0.05$) downregulated in tumor tissues are shown in red.

(C) Volcano plot of the CpG loci identified as cancer-specifically methylated in lung adenocarcinoma (normal $n = 4$, cancer $n = 19$) from TCGA Data Portal. The β value difference in DNA methylation between the tumor samples and the correspondent normal samples is plotted on the x axis, and the p value for a FDR-corrected Wilcoxon rank sum test of differences between the tumor and correspondent normal samples ($-1 \times \log_{10}$ scale) is plotted on the y axis. Probes that are significantly hypermethylated (FDR adjusted $p < 0.05$) in tumors are shown in red.

(D) Volcano plot gene expression data of cancer-specific DNA-methylated genes. Gene expression data were obtained from GEO (GSE7670) from primary lung adenocarcinoma ($n = 27$) and primary lung ($n = 30$). For the volcano plots, gene expression fold change between the normal tissues and the tumor tissues is plotted on the x axis, and the p value for a FDR-corrected t test of differences between the normal and the tumor tissues ($-1 \times \log_{10}$ scale) is plotted on the y axis. Probes that are significantly ($p < 0.05$) downregulated in tumor tissues are shown in red.

(E) Venn diagram showing the overlap between the genes statistically hypermethylated in colon adenocarcinoma ($n = 50$; FDR adjusted $p < 0.05$) and lung adenocarcinoma ($n = 33$; FDR adjusted $p < 0.05$).

(F) Venn diagram showing the overlap between the genes statistically repressed in colon adenocarcinoma ($n = 44$; FDR adjusted $p < 0.05$) and lung adenocarcinoma ($n = 25$; FDR adjusted $p < 0.05$).




Figure 4. Apoptosis Analysis of HCT116 and DKO1 Cells

(A) HCT116 wild-type and HCT116 DKO1 cells were stained with annexin V-FITC and PI and analyzed by FACS, showing an increased level of basal apoptotic cell death in the HCT116 DKO1 cell line compared to HCT116 wild-type. HCT116 DKO1 cells were then sorted in viable (annexin V and PI negative) and early apoptosis (annexin V positive and PI negative).

(B) The morphology of viable DKO1 and early apoptosis is clearly distinct. The apoptotic cells (blue) show a characteristic phenotype of higher SSC and lower FSC than the viable cells (red).

To gain more detailed information of how cancer cells become dependent on DNA methylation of these genes, we investigated how the silencing of one candidate, interleukin-1 receptor-associated kinase 3 (*IRAK3*), affects cancer cell survival in more detail. *IRAK3* has a cancer-specific DNA methylation pattern, a reduced expression in colon adenocarcinoma compared to normal colon, and a decreased DNA methylation in spontaneously dying DKO1 cells when compared to viable DKO1 cells. In addition, *IRAK3* was a promising candidate because, through *IRAK1* (Kobayashi et al., 2002), it indirectly inhibits three essential pathways that cancer cells rely on to survive: STAT3, NFkB, and MAPK (Figure S3F) (Ngo et al., 2011; Su et al., 2009; Turnis et al., 2010). These pathways, in turn, regulate the expression of the antiapoptotic gene *SURVIVIN* (Jiang et al., 2011; Zhou et al., 2009). Consistent with our hypothesis, using the Oncomine platform (<http://www.oncomine.org>), we found reduced *IRAK3* expression in several types of cancer when compared to normal tissue (Figure 5C) including colon adenocarcinoma when compared to normal colon (Student's t test $p = 6.59\text{E-}4$, top 15% underexpressed gene rank) (Kaiser et al., 2007), in lung adenocarcinoma when compared to normal lung (Student's t test $p = 5.70\text{E-}7$, top 6% underexpressed gene rank) (Su et al., 2007), in prostate carcinoma when compared to normal prostate (Student's t test $p = 4.01\text{E-}5$, top 3% underexpressed gene rank) (Welsh et al., 2001), and in cutaneous melanoma when compared to normal skin (Student's t test $p = 3.74\text{E-}8$, top 3% underexpressed gene rank) (Talantov et al., 2005) (Figure 5C). The consistently reduced expression level of *IRAK3* in a variety of cancers suggests that its silencing plays a role in the tumorigenic process. In addition the downregulation of *IRAK3* was correlated with a statistically significant upregulation of *IRAK1* and *SURVIVIN* in the same studies (Figures S3C and S3D).

To formally test whether the decreased expression of *IRAK3* was directly responsible for the increased expression of *SURVIVIN* and, consequently, increased cell survival, we re-expressed *IRAK3* in HCT116 cells. Re-expression of *IRAK3* caused a striking reduction in Survivin protein levels (Figure 5D) and caused a significant increase in cell death (Student's t test $p = 0.0219$; Figure 5E) and decrease in cell viability ($p < 0.0001$ Figure 5E), confirming that cancer cells require DNA methylation-induced silencing of *IRAK3*, and thus become dependent on the aberrant DNA methylation. Moreover, we sought to determine whether the silencing of *IRAK3* has any effect in a nontransformed cell. We performed a colony formation assay in a nontumorigenic cell line, URO-TSA, infected with *IRAK3* shRNA or a scrambled shRNA (Figure 5F). We observed that *IRAK3* knockdown was sufficient to induce a striking increase in colony formation (Figure 5F), thus demonstrating that the criteria we used were successful to identify functionally relevant genes and demonstrate that cancer cells become addicted to their epigenetic silencing.

DISCUSSION

Several genome-wide studies have revealed that a large number of promoter regions become de novo methylated in cancer (Noushmehr et al., 2010; Portela and Esteller, 2010). However, defining the specific “driver” gene regions that cancer cells depend on for survival has proven extremely difficult (Kalari and Pfeifer, 2010). In this study we have defined the gene promoters whose DNA methylation is required for survival of somatic cancer cells in culture. This group of genes could be further subdivided into at least three subgroups: those necessary to be methylated for the survival of (1) somatic cells, (2) cancer cells, and (3) cells in culture. These sets of genes retain DNA hypermethylation even after strong depletion of DNA methyltransferase activity, suggesting that DNA methylation is the main epigenetic mechanism used to maintain silencing because these cells do not seem able to switch to other repression mechanisms such as histone modifications alone.

Genes with germline-specific expression need to be tightly regulated in somatic tissues because their aberrant expression could be lethal for somatic cells. For example the gene “*Stil*,” in *Drosophila*, is only expressed in germ cells and is necessary for germ cell survival. When “*Stil*” is transiently expressed in somatic tissues, it results in lethality (Sahut-Barnola and Pauli, 1999). The genes we identified in the somatic-specific DNA methylation group are mainly germ cell-specific genes, and their demethylation and resulting re-expression in somatic cells can trigger apoptosis, as we showed for *SYCP3* and *ADAM2*. This suggests a primary role for DNA methylation as a mechanism for repression of testes-specific genes in somatic cells (which includes cancer cells). It also lends confidence to our analysis because it is known that several CpG island genes are normally DNA methylated in somatic tissue and unmethylated in germ cells (Shen et al., 2007).

Intriguingly, none of the genes identified in the cancer-specific group, whose DNA methylation is necessary for the survival of cancer cells, is classically known TSGs. This suggests that the genes we identified here are previously unknown tumor suppressors whose silencing is necessary for cancer cell survival. Interestingly, this group encompasses several cell signaling molecules, such as those with nucleotide receptor activity and G protein-coupled receptors (GPCRs). GPCRs have recently been described to be significantly mutated in several kinds of cancer and were found as a top category in a systematic search for TSGs by exome and transcriptome sequencing (Kan et al., 2010; Zhao et al., 2010). Taken together, these results indicate a more significant role of GPCRs, as TSGs, in cancer than previously thought.

TSGs can be silenced in cancer cells by several epigenetic mechanisms including DNA methylation, histone modifications, and nucleosome positioning (Jones and Baylin, 2007). It is also known that cancer cells depend on the silencing of TSGs and, consequently, are hypersensitive to the re-expression of these

(C) Bisulfite-sequencing analysis of CpG methylation status of four regions, from cancer-specific methylated cluster (*EYA4* and *IRAK3*) and from somatic tissue-specific DNA methylation cluster (*SYCP3* and *ADAM2*). The mean percent methylation at each CpG site is derived from clones shown on Figure S2A. The capped line represents the region analyzed.

See also Figure S2.




Figure 5. Functional Validations

(A) Overexpression of nine candidate genes from the cancer cluster (*P2RY14*, *IRAK3*, *CDO1*, *ESX1*, *ARMCX1*, *BCHE*), somatic cluster (*SYCP3* and *ADAM2*), and cell culture cluster (*STEAP4*). Shown is the fraction of Empty-Vector at the indicated times, normalized to the day 0 values. *NOX4* was used as a control gene because it is hypermethylated in HCT116 cells and completely demethylated in DKO1 cells. Data represent the mean \pm SEM.

(B) Overexpression of the same nine candidate genes reduces viability of RKO cancer cells. Shown is the fraction of Empty-Vector at the indicated times, normalized to the day 0 values. *NOX4* was used as a control gene. Data represent the mean \pm SEM.

genes (Luo et al., 2009; Martins et al., 2006; Ventura et al., 2007; Xue et al., 2007). We propose here that because of this TSG hypersensitivity, and because some of these key TSGs are silenced by epigenetic mechanisms, cancer cells become addicted to this aberrant epigenetic silencing.

Indeed, we demonstrated that re-expression of the genes we identified here decreases cell viability, highlighting the need for their constant repression. Furthermore, we showed that one of these genes, *IRAK3*, negatively regulates expression of the antiapoptotic gene *SURVIVIN*. It was recently shown that oncogenic activation of MYD88 led to IRAK1 phosphorylation and, consequently, NF κ B activation, promoting cell survival (Ngo et al., 2011). IRAK3 is a negative regulator of this signaling pathway, inhibiting IRAK1 phosphorylation (Janssens and Beyaert, 2003), thus supporting our findings and suggesting a tumor suppressor role for IRAK3. *IRAK3* was also previously identified, in an RNAi-based genetic screen, as able to suppress transformation of human mammary epithelial cells (Westbrook et al., 2005), consistent with our finding that *IRAK3* knockdown in nontumorigenic cells increased colony formation and suggesting that the genes found in this cancer group are excellent targets for therapy. In addition because these genes depend on DNA methylation-mediated gene silencing, these may be especially good targets for epigenetic therapy (Kelly et al., 2010).

It has been known for many years that de novo DNA methylation occurs during the cell-culturing process (Antequera et al., 1990; Jones et al., 1990; Wilson and Jones, 1983). Here, we identified a group of genes whose silencing by DNA methylation is required for cells to survive in culture. This group of genes was highly methylated in colon, bladder, and breast cell lines and unmethylated in primary-matched tissue analyzed, independent of the tumorigenic state (data not shown). Intriguingly, this group encompasses many nucleosome assembly genes, including several histone variants about which little is known. These results suggest that during the cell-culturing process, extensive changes in expression of nucleosome constituents are necessary for cell survival. It also highlights the importance of careful interpretation of epigenetic results obtained from cell culture experiments.

Taken together, by identifying the minimal DNA methylation profile necessary for the survival of cancer cells and comparing this profile in several primary normal tissues and cancer types, we were able to find, and experimentally validate, a group of genes whose de novo methylation in cancer is functionally relevant for the survival of cancer cells. We also found that, despite the complex nature of tumorigenesis, cancer cells

become dependent on the DNA methylation-mediated epigenetic silencing of these genes. These driver epigenetic events associated with cancer cell survival are potentially good candidates for the development of new, target-specific, therapies.

EXPERIMENTAL PROCEDURES

Cell Lines, DNA and RNA Preparations, Antibodies, and Primers

These are described in the Supplemental Experimental Procedures.

DNA Methylation Assay

Genomic DNA samples (1 μ g each) were bisulfite converted using the Zymo EZ DNA methylation kit (Zymo Research, Orange, CA, USA; catalog #D5002) according to the manufacturer's instructions. Bisulfite-converted DNA was eluted in an 18 μ l vol, and 3 μ l was removed for post-bisulfite quality control tests as described previously by Campan et al. (2009). All cell lines and clinical samples passed bisulfite conversion quality control and were subsequently processed for the Illumina Infinium DNA methylation platform (HumanMethylation27 BeadChip). A β value of 0–1.0 was reported for each CpG site (methylation from 0% to 100%, respectively). β Values were calculated as described previously by Wolff et al. (2010).

The Infinium methylation assays were performed by the USC Epigenome Center in accordance with the manufacturer's instructions. The assay information is available at <http://www.illumina.com>. Heat maps were generated for the β values. All the DNA methylation data from primary tissue was obtained from TCGA (<http://cancergenome.nih.gov/>).

To analyze the DNA methylation status of individual DNA molecules, we cloned bisulfite-converted PCR fragments into the pCR2.1 vector using the TOPO-TA cloning kit (Invitrogen, Carlsbad, CA, USA). Individual colonies were screened for the insert, and the region of interest was sequenced using M13 primers as previously described by Wolff et al. (2010).

H3K27me3 Status in ESCs as a Predictive Method for DNA Methylation in Cancer

H3K27me3 profile in H9 ESCs was obtained from previously published data by Lee et al. (2006). Next, we intersected the genomic position of the Infinium probes with the H3K27me3 status in H9 ESCs to define which probes are H3K27me3 positive and which are H3K27me3 negative in ESCs. From the approximately 24,000 probes used in Figure 1, we found approximately 7.7% as H3K27me3 positive in ESCs and 79.7% as H3K27me3 negative in ESCs, and we could not determine the H3K27me3 status in ESCs of 12.6% of the probes.

Genomic Architecture as a Predictive Method for DNA Methylation in Cancer

We used previously published data from Estéicio et al. (2010) that predict the inherent susceptibility to DNA methylation in cancer based on SINE and LINE retrotransposon density in a 20 kb window around the TSS of each gene. Briefly, they calculate the log odds ratio of SINE and LINE retrotransposons per 1 kb window, and the sum of log odds scores in the 20 kb region allowed the classification of each gene as methylation prone, methylation intermediate, and methylation resistant.

(C) Metaanalysis using the Oncomine (<http://www.oncomine.org>) for *IRAK3* expression. Box plots showing decreased expression of *IRAK3* during tumorigenesis on data sets performed in colon adenocarcinoma (Kaiser et al., 2007), lung adenocarcinoma (Su et al., 2007), prostate carcinoma (Welsh et al., 2001), and cutaneous melanoma (Talantov et al., 2005). The y axis represents log₂ median-centered intensity (normalized expression). Shaded boxes represent the interquartile range (25th–75th percentile). Whiskers represent the 10th–90th percentile. The bars denote the median.

(D) Overexpression of *IRAK3* in HCT116 cells induces a reduction in the Survivin levels. Western blot analyses of IRAK3 and Survivin after lentiviral infection with pLJM1 empty vector (EV) or pLJM1-IRAK3. Histone H3 was used as a loading control.

(E) *IRAK3* expression induces cell death of cancer cells. HCT116 infected with pLJM1 empty vector or pLJM1 *IRAK3* was stained with annexin V-FITC and PI and analyzed by FACS, showing an increased level of cell death in the cell overexpressing *IRAK3* (upper panel). Re-expression of *IRAK3* in HCT116 wild-type cells showed a reduced cell number in culture than HCT116 empty vector (lower panel). * $p < 0.05$, *** $p < 0.0001$. Data represent the mean \pm SEM.

(F) *IRAK3* knockdown induces colony formation in a nontumorigenic cell. UROTSa infected with a shRNA against *IRAK3* presented a higher colony formation activity than UROTSa infected with a scrambled shRNA. Western blot analysis of IRAK3 after lentiviral infection with shRNA against *IRAK3* or a scrambled shRNA was performed. Actin was used as a loading control.

See also Figure S3.

Apoptosis Assay

Cellular apoptosis was measured by annexin V and propidium iodide (PI) staining using annexin V-FITC Apoptosis Detection Kit (MBL), according to the manufacturer's protocol. Following staining the cells were analyzed and sorted by FACS analysis as described previously by De Carvalho et al. (2011).

Gene Expression Analysis

All gene expression data from primary tissue were obtained from GEO (GSE8671 and GSE7670). The data were median normalized and log2 transformed.

Ectopic Gene Expression

This is described in the [Supplemental Experimental Procedures](#).

Statistical Analysis

This is described in the [Supplemental Experimental Procedures](#).

ACCESSION NUMBERS

Infinium methylation data were deposited in NCBI's Gene Expression Omnibus: series accession number GSE36534.

SUPPLEMENTAL INFORMATION

Supplemental Information includes three figures, one table, and Supplemental Experimental Procedures and can be found with this article online at [doi:10.1016/j.ccr.2012.03.045](https://doi.org/10.1016/j.ccr.2012.03.045).

ACKNOWLEDGMENTS

This work was supported by R37CA082422 (to P.A.J.). We would like to thank Drs. Bert Vogelstein and Stephen B. Baylin for kindly providing DKO cells and also Lora Barsky from the Broad CIRM Center, Flow Cytometry Core Facility for technical assistance with FACS analysis. We thank the members of the P.A.J. laboratory and Daniel Weisenberger for helpful discussions and for careful reading of the manuscript. We thank Kimberly Siegmund for the helpful discussions about our statistical approaches. We also thank Peter Laird for useful discussions and The Cancer Genome Atlas.

Received: February 26, 2011

Revised: August 8, 2011

Accepted: March 27, 2012

Published: May 14, 2012

REFERENCES

- Antequera, F., Boyes, J., and Bird, A. (1990). High levels of de novo methylation and altered chromatin structure at CpG islands in cell lines. *Cell* 62, 503–514.
- Baylin, S.B., and Ohm, J.E. (2006). Epigenetic gene silencing in cancer—a mechanism for early oncogenic pathway addiction? *Nat. Rev. Cancer* 6, 107–116.
- Campan, M., Weisenberger, D.J., Trinh, B., and Laird, P.W. (2009). MethyLight. *Methods Mol. Biol.* 507, 325–337.
- Cedar, H., and Bergman, Y. (2009). Linking DNA methylation and histone modification: patterns and paradigms. *Nat. Rev. Genet.* 10, 295–304.
- Chen, T., Hevi, S., Gay, F., Tsujimoto, N., He, T., Zhang, B., Ueda, Y., and Li, E. (2007). Complete inactivation of DNMT1 leads to mitotic catastrophe in human cancer cells. *Nat. Genet.* 39, 391–396.
- Darzynkiewicz, Z., Bruno, S., Del Bino, G., Gorczyca, W., Hotz, M.A., Lassota, P., and Traganos, F. (1992). Features of apoptotic cells measured by flow cytometry. *Cytometry* 13, 795–808.
- De Carvalho, D.D., You, J.S., and Jones, P.A. (2010). DNA methylation and cellular reprogramming. *Trends Cell Biol.* 20, 609–617.
- De Carvalho, D.D., Binato, R., Pereira, W.O., Leroy, J.M., Colassanti, M.D., Proto-Siqueira, R., Bueno-Da-Silva, A.E., Zago, M.A., Zanichelli, M.A., Abdelhay, E., et al. (2011). BCR-ABL-mediated upregulation of PRAME is responsible for knocking down TRAIL in CML patients. *Oncogene* 30, 223–233.
- Del Prete, M.J., Robles, M.S., Guão, A., Martínez-A, C., Izquierdo, M., and Garcia-Sanz, J.A. (2002). Degradation of cellular mRNA is a general early apoptosis-induced event. *FASEB J.* 16, 2003–2005.
- Egger, G., Jeong, S., Escobar, S.G., Cortez, C.C., Li, T.W., Saito, Y., Yoo, C.B., Jones, P.A., and Liang, G. (2006). Identification of DNMT1 (DNA methyltransferase 1) hypomorphs in somatic knockouts suggests an essential role for DNMT1 in cell survival. *Proc. Natl. Acad. Sci. USA* 103, 14080–14085.
- Estécio, M.R., Gallegos, J., Vallot, C., Castoro, R.J., Chung, W., Maegawa, S., Oki, Y., Kondo, Y., Jelinek, J., Shen, L., et al. (2010). Genome architecture marked by retrotransposons modulates predisposition to DNA methylation in cancer. *Genome Res.* 20, 1369–1382.
- Figuroa, M.E., Lugthart, S., Li, Y., Erpelinck-Verschueren, C., Deng, X., Christos, P.J., Schifano, E., Booth, J., van Putten, W., Skrabanek, L., et al. (2010). DNA methylation signatures identify biologically distinct subtypes in acute myeloid leukemia. *Cancer Cell* 17, 13–27.
- Gal-Yam, E.N., Egger, G., Iniguez, L., Holster, H., Einarsson, S., Zhang, X., Lin, J.C., Liang, G., Jones, P.A., and Tanay, A. (2008). Frequent switching of Polycomb repressive marks and DNA hypermethylation in the PC3 prostate cancer cell line. *Proc. Natl. Acad. Sci. USA* 105, 12979–12984.
- Hanahan, D., and Weinberg, R.A. (2000). The hallmarks of cancer. *Cell* 100, 57–70.
- Hanahan, D., and Weinberg, R.A. (2011). Hallmarks of cancer: the next generation. *Cell* 144, 646–674.
- Irizarry, R.A., Ladd-Acosta, C., Wen, B., Wu, Z., Montano, C., Onyango, P., Cui, H., Gabo, K., Rongione, M., Webster, M., et al. (2009). The human colon cancer methylome shows similar hypo- and hypermethylation at conserved tissue-specific CpG island shores. *Nat. Genet.* 41, 178–186.
- Janssens, S., and Beyaert, R. (2003). Functional diversity and regulation of different interleukin-1 receptor-associated kinase (IRAK) family members. *Mol. Cell* 11, 293–302.
- Jiang, Y.F., Sr., He, B., Li, N.P., Ma, J., Gong, G.Z., and Zhang, M. (2011). The oncogenic role of NS5A of hepatitis C virus is mediated by up-regulation of survivin gene expression in the hepatocellular cell through p53 and NF- κ B pathways. *Cell Biol. Int.* 35, 1225–1232.
- Jones, P.A., and Baylin, S.B. (2002). The fundamental role of epigenetic events in cancer. *Nat. Rev. Genet.* 3, 415–428.
- Jones, P.A., and Baylin, S.B. (2007). The epigenomics of cancer. *Cell* 128, 683–692.
- Jones, P.A., Wolkowicz, M.J., Rideout, W.M., 3rd, Gonzales, F.A., Marziasz, C.M., Coetzee, G.A., and Tapscott, S.J. (1990). De novo methylation of the MyoD1 CpG island during the establishment of immortal cell lines. *Proc. Natl. Acad. Sci. USA* 87, 6117–6121.
- Kaiser, S., Park, Y.K., Franklin, J.L., Halberg, R.B., Yu, M., Jessen, W.J., Freudenberger, J., Chen, X., Haigis, K., Jegga, A.G., et al. (2007). Transcriptional recapitulation and subversion of embryonic colon development by mouse colon tumor models and human colon cancer. *Genome Biol.* 8, R131.
- Kalari, S., and Pfeifer, G.P. (2010). Identification of driver and passenger DNA methylation in cancer by epigenomic analysis. *Adv. Genet.* 70, 277–308.
- Kan, Z., Jaiswal, B.S., Stinson, J., Janakiraman, V., Bhatt, D., Stern, H.M., Yue, P., Haverly, P.M., Bourgon, R., Zheng, J., et al. (2010). Diverse somatic mutation patterns and pathway alterations in human cancers. *Nature* 466, 869–873.
- Kelly, T.K., De Carvalho, D.D., and Jones, P.A. (2010). Epigenetic modifications as therapeutic targets. *Nat. Biotechnol.* 28, 1069–1078.
- Kobayashi, K., Hernandez, L.D., Galán, J.E., Janeway, C.A., Jr., Medzhitov, R., and Flavell, R.A. (2002). IRAK-M is a negative regulator of Toll-like receptor signaling. *Cell* 110, 191–202.
- Lee, T.I., Jenner, R.G., Boyer, L.A., Guenther, M.G., Levine, S.S., Kumar, R.M., Chevalier, B., Johnstone, S.E., Cole, M.F., Isono, K., et al. (2006). Control of developmental regulators by Polycomb in human embryonic stem cells. *Cell* 125, 301–313.

- Luo, J., Solimini, N.L., and Elledge, S.J. (2009). Principles of cancer therapy: oncogene and non-oncogene addiction. *Cell* 136, 823–837.
- Manns, J., Daubrawa, M., Driessen, S., Paasch, F., Hoffmann, N., Löffler, A., Lauber, K., Dieterle, A., Alers, S., Iftner, T., et al. (2011). Triggering of a novel intrinsic apoptosis pathway by the kinase inhibitor staurosporine: activation of caspase-9 in the absence of Apaf-1. *FASEB J.* 25, 3250–3261.
- Martins, C.P., Brown-Swigart, L., and Evan, G.I. (2006). Modeling the therapeutic efficacy of p53 restoration in tumors. *Cell* 127, 1323–1334.
- Meissner, A. (2010). Epigenetic modifications in pluripotent and differentiated cells. *Nat. Biotechnol.* 28, 1079–1088.
- Ngo, V.N., Young, R.M., Schmitz, R., Jhavar, S., Xiao, W., Lim, K.H., Kohlhammer, H., Xu, W., Yang, Y., Zhao, H., et al. (2011). Oncogenically active MYD88 mutations in human lymphoma. *Nature* 470, 115–119.
- Noushmehr, H., Weisenberger, D.J., Diefes, K., Phillips, H.S., Pujara, K., Berman, B.P., Pan, F., Pelloski, C.E., Sulman, E.P., Bhat, K.P., et al; Cancer Genome Atlas Research Network. (2010). Identification of a CpG island methylator phenotype that defines a distinct subgroup of glioma. *Cancer Cell* 17, 510–522.
- Ohm, J.E., McGarvey, K.M., Yu, X., Cheng, L., Schuebel, K.E., Cope, L., Mohammad, H.P., Chen, W., Daniel, V.C., Yu, W., et al. (2007). A stem cell-like chromatin pattern may predispose tumor suppressor genes to DNA hypermethylation and heritable silencing. *Nat. Genet.* 39, 237–242.
- Portela, A., and Esteller, M. (2010). Epigenetic modifications and human disease. *Nat. Biotechnol.* 28, 1057–1068.
- Rhee, I., Bachman, K.E., Park, B.H., Jair, K.W., Yen, R.W., Schuebel, K.E., Cui, H., Feinberg, A.P., Lengauer, C., Kinzler, K.W., et al. (2002). DNMT1 and DNMT3b cooperate to silence genes in human cancer cells. *Nature* 416, 552–556.
- Robertson, K.D. (2005). DNA methylation and human disease. *Nat. Rev. Genet.* 6, 597–610.
- Sabates-Bellver, J., Van der Flier, L.G., de Palo, M., Cattaneo, E., Maake, C., Rehrauer, H., Laczko, E., Kurowski, M.A., Bujnicki, J.M., Menigatti, M., et al. (2007). Transcriptome profile of human colorectal adenomas. *Mol. Cancer Res.* 5, 1263–1275.
- Sahut-Barnola, I., and Pauli, D. (1999). The *Drosophila* gene stand still encodes a germline chromatin-associated protein that controls the transcription of the ovarian tumor gene. *Development* 126, 1917–1926.
- Schlesinger, Y., Straussman, R., Keshet, I., Farkash, S., Hecht, M., Zimmerman, J., Eden, E., Yakhini, Z., Ben-Shushan, E., Reubinoff, B.E., et al. (2007). Polycomb-mediated methylation on Lys27 of histone H3 pre-marks genes for de novo methylation in cancer. *Nat. Genet.* 39, 232–236.
- Sharma, S., Kelly, T.K., and Jones, P.A. (2010). Epigenetics in cancer. *Carcinogenesis* 31, 27–36.
- Sharma, S., De Carvalho, D.D., Jeong, S., Jones, P.A., and Liang, G. (2011). Nucleosomes containing methylated DNA stabilize DNA methyltransferases 3A/3B and ensure faithful epigenetic inheritance. *PLoS Genet.* 7, e1001286.
- Shen, L., Kondo, Y., Guo, Y., Zhang, J., Zhang, L., Ahmed, S., Shu, J., Chen, X., Waterland, R.A., and Issa, J.P. (2007). Genome-wide profiling of DNA methylation reveals a class of normally methylated CpG island promoters. *PLoS Genet.* 3, 2023–2036.
- Solimini, N.L., Luo, J., and Elledge, S.J. (2007). Non-oncogene addiction and the stress phenotype of cancer cells. *Cell* 130, 986–988.
- Spada, F., Haemmer, A., Kuch, D., Rothbauer, U., Schermelleh, L., Kremmer, E., Carell, T., Längst, G., and Leonhardt, H. (2007). DNMT1 but not its interaction with the replication machinery is required for maintenance of DNA methylation in human cells. *J. Cell Biol.* 176, 565–571.
- Su, J., Zhang, T., Tyson, J., and Li, L. (2009). The interleukin-1 receptor-associated kinase M selectively inhibits the alternative, instead of the classical NFκB pathway. *J. Innate Immun.* 1, 164–174.
- Su, L.J., Chang, C.W., Wu, Y.C., Chen, K.C., Lin, C.J., Liang, S.C., Lin, C.H., Whang-Peng, J., Hsu, S.L., Chen, C.H., and Huang, C.Y. (2007). Selection of DDX5 as a novel internal control for Q-RT-PCR from microarray data using a block bootstrap re-sampling scheme. *BMC Genomics* 8, 140.
- Taberlay, P.C., and Jones, P.A. (2011). DNA methylation and cancer. *Prog. Drug Res.* 67, 1–23.
- Talantov, D., Mazumder, A., Yu, J.X., Briggs, T., Jiang, Y., Backus, J., Atkins, D., and Wang, Y. (2005). Novel genes associated with malignant melanoma but not benign melanocytic lesions. *Clin. Cancer Res.* 11, 7234–7242.
- Turnis, M.E., Song, X.T., Bear, A., Foster, A.E., Gottschalk, S., Brenner, M.K., Chen, S.Y., and Rooney, C.M. (2010). IRAK-M removal counteracts dendritic cell vaccine deficits in migration and longevity. *J. Immunol.* 185, 4223–4232.
- Ventura, A., Kirsch, D.G., McLaughlin, M.E., Tuveson, D.A., Grimm, J., Lintault, L., Newman, J., Reczek, E.E., Weissleder, R., and Jacks, T. (2007). Restoration of p53 function leads to tumour regression in vivo. *Nature* 445, 661–665.
- Weinstein, I.B. (2002). Cancer. Addiction to oncogenes—the Achilles heel of cancer. *Science* 297, 63–64.
- Welsh, J.B., Sapinoso, L.M., Su, A.I., Kern, S.G., Wang-Rodriguez, J., Moskaluk, C.A., Frierson, H.F., Jr., and Hampton, G.M. (2001). Analysis of gene expression identifies candidate markers and pharmacological targets in prostate cancer. *Cancer Res.* 61, 5974–5978.
- Westbrook, T.F., Martin, E.S., Schlabach, M.R., Leng, Y., Liang, A.C., Feng, B., Zhao, J.J., Roberts, T.M., Mandel, G., Hannon, G.J., et al. (2005). A genetic screen for candidate tumor suppressors identifies REST. *Cell* 121, 837–848.
- Widschwendter, M., Fiegl, H., Egle, D., Mueller-Holzner, E., Spizzo, G., Marth, C., Weisenberger, D.J., Campan, M., Young, J., Jacobs, I., and Laird, P.W. (2007). Epigenetic stem cell signature in cancer. *Nat. Genet.* 39, 157–158.
- Wilson, V.L., and Jones, P.A. (1983). DNA methylation decreases in aging but not in immortal cells. *Science* 220, 1055–1057.
- Wolff, E.M., Chihara, Y., Pan, F., Weisenberger, D.J., Siegmund, K.D., Sugano, K., Kawashima, K., Laird, P.W., Jones, P.A., and Liang, G. (2010). Unique DNA methylation patterns distinguish noninvasive and invasive urothelial cancers and establish an epigenetic field defect in premalignant tissue. *Cancer Res.* 70, 8169–8178.
- Xue, W., Zender, L., Miething, C., Dickins, R.A., Hernando, E., Krizhanovskiy, V., Cordon-Cardo, C., and Lowe, S.W. (2007). Senescence and tumour clearance is triggered by p53 restoration in murine liver carcinomas. *Nature* 445, 656–660.
- Zhao, Q., Kirkness, E.F., Caballero, O.L., Galante, P.A., Parmigiani, R.B., Edsall, L., Kuan, S., Ye, Z., Levy, S., Vasconcelos, A.T., et al. (2010). Systematic detection of putative tumor suppressor genes through the combined use of exome and transcriptome sequencing. *Genome Biol.* 11, R114.
- Zhou, J., Bi, C., Janakakumara, J.V., Liu, S.C., Chng, W.J., Tay, K.G., Poon, L.F., Xie, Z., Palaniyandi, S., Yu, H., et al. (2009). Enhanced activation of STAT pathways and overexpression of survivin confer resistance to FLT3 inhibitors and could be therapeutic targets in AML. *Blood* 113, 4052–4062.